

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 October 2001 (18.10.2001)

PCT

(10) International Publication Number
WO 01/76601 A2

- (51) International Patent Classification⁷: **A61K 31/46**, 31/137 // (A61K 31/46, 31:137)
- (21) International Application Number: **PCT/EP01/03889**
- (22) International Filing Date: **5 April 2001 (05.04.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
0008485.5 **7 April 2000 (07.04.2000)** **GB**
- (71) Applicant (for all designated States except US): **GLAXO GROUP LIMITED [GB/GB]**; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **JENKINS, Richard, John [GB/US]**; Glaxo Wellcome Inc., Five Moore Drive, Research Triangle Park, NC 27709 (US). **RICHARDS, David, Hugh [GB/GB]**; Glaxo Wellcome Plc, Stockley Park West, Uxbridge, Middlesex UB11 1BT (GB).
- (74) Agent: **LEAROYD, Stephanie, Anne**; Glaxo Smith Kline, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).
- (81) Designated States (national): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.**
- (84) Designated States (regional): **ARIPO** patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), **Eurasian** patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), **European** patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), **OAPI** patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/76601 A2

(54) Title: **PHARMACEUTICAL COMPOSITIONS**

(57) Abstract: The present invention is concerned with pharmaceutical formulations comprising a combination of salmeterol and ipratropium and the use of such formulations in medicine, particularly in the prophylaxis and treatment of respiratory diseases.

Pharmaceutical Compositions

The present invention is concerned with compositions containing a combination of salmeterol and ipratropium and the use of such compositions in medicine,
5 particularly in the prophylaxis and treatment of respiratory diseases.

GB 2 140 800 describes phenethanolamine compounds which are β_2 -adrenoreceptor agonists including 4-hydroxy- α^1 -[[[6-(4-phenylbutoxy)hexyl]-amino]methyl]-1,3-benzenedimethanol 1-hydroxy-2-naphthalenecarboxylate
10 (salmeterol xinafoate) which is now used clinically in the treatment of bronchial asthma and related disorders.

US 3,505,337 and US 3,681,500 describe ipratropium and its salts, such (endo,syn)-(+/-)-3-(3-hydroxy-1-oxo-2-phenylpropoxy)-8-methyl-8-(1-methylethyl)-8-azoniabicyclo[3.2.1]octane bromide (ipratropium bromide) and
15 pharmaceutical formulations thereof. Ipratropium bromide is an anticholinergic agent, which is now used clinically in the treatment of bronchial asthma and related disorders.

20 Although salmeterol xinafoate and ipratropium bromide are effective bronchodilators, the maximum duration of action is 12 hours and 6 hours respectively, hence twice daily or four times daily dosing is often required. There is therefore a clinical need for bronchodilators having potent and selective action and having an advantageous profile of action

25 According to the present invention, there is provided a pharmaceutical formulation comprising salmeterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and ipratropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or excipient, and optionally
30 one or more other therapeutic ingredients. According to a preferred aspect of the present invention, there is provided a pharmaceutical formulation comprising salmeterol xinafoate and ipratropium bromide, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic

ingredients. In the most preferred aspect, the above pharmaceutical formulations are suitable for administration by inhalation.

5 It is to be understood that the present invention covers all combinations of particular and preferred aspects of the invention described herein.

As would be appreciated by the skilled person, salmeterol includes an asymmetric centre and ipratropium includes three asymmetric centres. The present invention includes both (S) and (R) enantiomers of salmeterol either in substantially pure form or admixed in any proportions, as well as each isomer of
10 ipratropium either in substantially pure form or admixed in any proportions. The enantiomers of salmeterol have been described previously, for example, in EP0422889 and WO 99/13867. Various isomers of ipratropium are described in DE 4140861 and WO 97/05136.

15 By the term "physiologically functional derivative" is meant a chemical derivative of salmeterol or ipratropium having the same physiological function as the free compound, for example, by being convertible in the body thereto. According to the present invention, examples of physiologically functional derivatives include esters.

20 Suitable salts according to the invention include those formed with both organic and inorganic acids. Pharmaceutically acceptable acid addition salts include but are not limited to those formed from hydrochloric, hydrobromic, sulphuric, citric, tartaric, phosphoric, lactic, pyruvic, acetic, trifluoroacetic, succinic, oxalic,
25 fumaric, maleic, oxaloacetic, methanesulphonic, ethanesulphonic, p-toluenesulphonic, benzenesulphonic, isethionic, and naphthalenecarboxylic, such as 1-hydroxy-2-naphthalenecarboxylic acids.

30 Pharmaceutically acceptable esters of salmeterol or ipratropium may have a hydroxyl group converted to a C₁₋₆alkyl, aryl, aryl C₁₋₆ alkyl, or amino acid ester.

As mentioned above, both salmeterol and ipratropium and their pharmaceutically acceptable salts, solvates, and physiologically functional derivatives have been described for use in the treatment of respiratory diseases. Therefore,

formulations of salmeterol and ipratropium and their pharmaceutically acceptable salts, solvates, and physiologically functional derivatives have use in the prophylaxis and treatment of clinical conditions for which a selective β_2 -adrenoreceptor agonist and/or an anticholinergic agent is indicated. Such

5 conditions include diseases associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary diseases (COPD) (e.g. chronic and wheezy bronchitis, emphysema), respiratory tract infection and upper respiratory tract disease (e.g. rhinitis, such as allergic and seasonal rhinitis).

10 Accordingly, the present invention provides a method for the prophylaxis or treatment of a clinical condition in a mammal, such as a human, for which a selective β_2 -adrenoreceptor agonist and/or anticholinergic agent is indicated, which comprises administration of a therapeutically effective amount of a pharmaceutical formulation comprising salmeterol or a pharmaceutically

15 acceptable salt, solvate, or physiologically functional derivative thereof and ipratropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or excipient. In a preferred aspect, there is provided such a method which comprises administration of a therapeutically effective amount of a

20 pharmaceutical formulation comprising salmeterol xinafoate and ipratropium bromide, and a pharmaceutically acceptable carrier or excipient. In particular, the present invention provides such a method for the prophylaxis or treatment of a disease associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary disease (COPD), respiratory tract infection or

25 upper respiratory tract disease.

The amount of salmeterol and ipratropium, or a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof which is required to achieve a therapeutic effect will, of course, vary with the particular compound,

30 the route of administration, the subject under treatment, and the particular disorder or disease being treated. As a monotherapy, salmeterol xinafoate is generally administered to adult humans by aerosol inhalation at a dose of 50mcg or 100mcg twice daily. As a monotherapy, ipratropium bromide is generally administered to adult humans by inhalation at a dose of from 20mcg to 80mcg

35 three or four times daily.

Hereinafter, the term "active ingredients" means salmeterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, preferably salmeterol xinafoate, and ipratropium, or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, preferably ipratropium bromide.

Suitably, the pharmaceutical formulations which are suitable for inhalation according to the invention comprise the active ingredients in amounts such that each actuation provides therapeutically effective dose, for example, a dose of salmeterol of 10mcg to 150mcg, preferably 50mcg and a dose of ipratropium bromide of 10mcg to 400mcg, preferably 50mcg to 320mcg, more preferably, 80mcg to 160mcg. The pharmaceutical formulations according to the invention may further include other therapeutic agents for example anti-inflammatory agents such as corticosteroids or other β_2 -adrenoreceptor agonists (such as salbutamol, formoterol, fenoterol or terbutaline and salts thereof).

Suitably, the pharmaceutical formulations which are suitable for inhalation according to the invention provide therapeutically effective doses that permit the establishment of a twice daily (bis in diem – b.i.d) dosing regimen.

The formulations include those suitable for oral, parenteral (including subcutaneous, intradermal, intramuscular, intravenous and intraarticular), intranasal, inhalation (including fine particle dusts or mists which may be generated by means of various types of metered dose pressurised aerosols, nebulisers or insufflators), rectal and topical (including dermal, buccal, sublingual and intraocular) administration although the most suitable route may depend upon for example the condition and disorder of the recipient. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active ingredients into association with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

Formulations for inhalation include powder compositions which will preferably contain lactose, and spray compositions which may be formulated, for example, as aqueous solutions or suspensions or as aerosols delivered from pressurised packs, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, 1,1,1,2,3,3,3-heptafluoropropane, 1,1,1,2-tetrafluoroethane, carbon dioxide or other suitable gas. Suitable aerosol formulations include those described in EP 0372777 and WO93/11743. Suitable aerosol formulations may include excipients such as surfactants and/or cosolvents such as ethanol. For suspension aerosols, the active ingredients should be micronised so as to permit inhalation of substantially all of the active ingredients into the lungs upon administration of the aerosol formulation, thus the active ingredients will have a particle size of less than 100 microns, desirably less than 20 microns, and preferably in the range 1 to 10 microns, for example, 1 to 5 microns.

Preferred compositions for aerosol delivery consist essentially of particulate active ingredients, and 1,1,1,2-tetrafluoroethane, 1,1,1,2,3,3,3-heptafluoropropane or mixtures thereof as propellant. Especially preferred compositions for aerosol delivery consist of particulate active ingredients, and 1,1,1,2-tetrafluoroethane, 1,1,1,2,3,3,3-heptafluoropropane or mixtures thereof as propellant.

Compositions according to the present invention for aerosol delivery may be filled into canisters suitable for delivering pharmaceutical aerosol formulations. Canisters generally comprise a container capable of withstanding the vapour pressure of the propellant, such as plastic or plastic-coated glass bottle or preferably a metal can, for example an aluminium can which may optionally be anodised, lacquer-coated and/or plastic-coated, which container is closed with a metering valve. It may be preferred that canisters be coated with a fluorocarbon polymer as described in WO 96/32150, for example, a co-polymer of polyethersulphone (PES) and polytetrafluoroethylene (PTFE). Another polymer for coating that may be contemplated is FEP (fluorinated ethylene propylene). The metering valves are designed to deliver a metered amount of the formulation per actuation and incorporate a gasket to prevent leakage of

propellant through the valve. The gasket may comprise any suitable elastomeric material such as for example low density polyethylene, chlorobutyl, black and white butadiene-acrylonitrile rubbers, butyl rubber and neoprene. Thermoplastic elastomer valves as described in WO92/11190 and valves containing EPDM rubber as described in WO95/02650 are especially suitable. Suitable valves are commercially available from manufacturers well known in the aerosol industry, for example, from Valois, France (eg. DF10, DF30, DF60), Bepak plc, UK (eg. BK300, BK356, BK357) and 3M-Neotechnic Ltd, UK (eg. Spraymiser™). The DF31 valve of Valois, France is also suitable.

Valve seals, especially the gasket seal and also the seals around the metering chamber, will preferably be manufactured of a material which is inert to and resists extraction into the contents of the formulation, especially when the contents include ethanol.

Valve materials, especially the material of manufacture of the metering chamber, will preferably be manufactured of a material which is inert to and resists distortion by contents of the formulation, especially when the contents include ethanol. Particularly suitable materials for use in manufacture of the metering chamber include polyesters eg polybutyleneterephthalate (PBT) and acetals, especially PBT.

Materials of manufacture of the metering chamber and/or the valve stem may desirably be fluorinated, partially fluorinated or impregnated with fluorine containing substances in order to resist drug deposition.

Valves which are entirely or substantially composed of metal components (eg Spraymiser, 3M-Neotechnic) are especially preferred for use according to the invention.

Intranasal sprays may be formulated with aqueous or non-aqueous vehicles with the addition of agents such as thickening agents, buffer salts or acid or alkali to adjust the pH, isotonicity adjusting agents or anti-oxidants.

Capsules and cartridges or for example gelatin, or blisters of for example laminated aluminium foil, for use in an inhaler or insufflator may be formulated containing a powder mix of the active ingredients and a suitable powder base such as lactose or starch, preferably lactose. In this aspect, the active ingredients are suitably micronised so as to permit inhalation of substantially all of the active ingredients into the lungs upon administration of the dry powder formulation, thus the active ingredients will have a particle size of less than 100 microns, desirably less than 20 microns, and preferably in the range 1 to 10 microns.

Solutions for inhalation by nebulation may be formulated with an aqueous vehicle with the addition of agents such as acid or alkali, buffer salts, isotonicity adjusting agents or antimicrobials. They may be sterilised by filtration or heating in an autoclave, or presented as a non-sterile product.

Preferred unit dosage formulations are those containing a pharmaceutically effective dose, as hereinbefore recited, or an appropriate fraction thereof, of the active ingredient. Thus, in the case of formulations designed for delivery by metered dose pressurised aerosols, one actuation of the aerosol may deliver half of the therapeutically effective amount such that two actuations are necessary to deliver the therapeutically effective dose.

It should be understood that in addition to the ingredients particularly mentioned above, the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question. Furthermore, the claimed formulations include bioequivalents as defined by the US Food and Drugs Agency.

For a better understanding of the invention, the following Examples are given by way of illustration.

EXAMPLES

A: Metered Dose Inhalers

Example 1: 25/40 salmeterol/ipratropium

	Per Inhaler %w/w	Per actuation
Salmeterol Xinafoate	0.05	36.3 microgram
Ipratropium Bromide	0.05	40 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.0mg

5 The micronised active ingredients are weighed into an aluminium can, 1,1,1,2-tetrafluoroethane is then added from a vacuum flask and a metering valve with a 63 μ l metering chamber is crimped into place.

Similar methods may be used for the formulations of Examples 2 to 8:

Example 2: 25/60 salmeterol/ipratropium

10

	Per Inhaler %w/w	Per actuation
Salmeterol Xinafoate	0.05	36.3 microgram
Ipratropium Bromide	0.08	60 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.0mg

Example 3: 25/80 salmeterol/ipratropium

	Per Inhaler %w/w	Per actuation
Salmeterol Xinafoate	0.05	36.3 microgram
Ipratropium Bromide	0.11	80 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.0mg

15 Example 4: 25/160 salmeterol/ipratropium

	Per Inhaler %w/w	Per actuation
Salmeterol Xinafoate	0.05	36.3 microgram
Ipratropium Bromide	0.21	160 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.0mg

In Examples 5 to 8, a metering valve with a 50 μ l metering chamber is used.

Example 5: 25/40 salmeterol/ipratropium

	Per Inhaler %w/w	Per actuation
Salmeterol Xinafoate	0.06	36.3 microgram
Ipratropium Bromide	0.07	40 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 60.0mg

5 Example 6: 25/60 salmeterol/ipratropium

	Per Inhaler %w/w	Per actuation
Salmeterol Xinafoate	0.06	36.3 microgram
Ipratropium Bromide	0.10	60 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 60.0mg

Example 7: 25/80 salmeterol/ipratropium

	Per Inhaler %w/w	Per actuation
Salmeterol Xinafoate	0.06	36.3 microgram
Ipratropium Bromide	0.13	80 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 60.0mg

10

Example 8: 25/160 salmeterol/ipratropium

	Per Inhaler %w/w	Per actuation
Salmeterol Xinafoate	0.06	36.3 microgram
Ipratropium Bromide	0.27	160 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 60.0mg

Example 9: 25/80 salmeterol/ipratropium

15

	Per Inhaler %w/w	Per actuation
Salmeterol Xinafoate	0.05	36.3 microgram
Ipratropium Bromide	0.11	80 microgram

1,1,1,2-Tetrafluoroethane	to 100	to 75.0mg
---------------------------	--------	-----------

The micronised active ingredients were weighed into an aluminium can coated internally with a PTFE/PES polymer blend as described in WO96/32150, 1,1,1,2-tetrafluoroethane was then added from a vacuum flask and a Valois DF60 metering valve (metering chamber volume 63 μ l) was crimped into place.

In an alternative process, the formulation of Example 9 was also made up by weighing the micronised active ingredients into a pressurised vessel and adding a portion of the 1,1,1,2-tetrafluoroethane to form a suspension. Aliquots of the suspension were filled, through the valve, into a number of aluminium cans coated internally with a PTFE/PES polymer blend as described in WO96/32150 and closed with Valois DF60 metering valves. Further 1,1,1,2-tetrafluoroethane (to 100%w/w) was then added to each can through the valve.

B: Dry Powder Inhalers

Example 10: 50/80 salmeterol/ipratropium

	Per cartridge or blister
Salmeterol Xinafoate	72.6 microgram
Ipratropium Bromide	80 microgram
Lactose Ph. Eur.	to 12.5mg or to 25.0mg

The active ingredients are micronised and bulk blended with the lactose in the proportions given above. The blend is filled into hard gelatin capsules or cartridges or in specifically constructed double foil blister packs to be administered by an inhaler such as a Rotahaler, Diskhaler, or Diskus inhaler (each of these being a Trademark of Glaxo Group Limited).

Similar methods may be used for the formulations of Examples 11 to 13:

Example 11: 50/120 salmeterol/ipratropium

11

	Per cartridge or blister
Salmeterol Xinafoate	72.6 microgram
Ipratropium Bromide	120 microgram
Lactose Ph. Eur.	to 12.5mg or to 25.0mg

Example 12: 50/160 salmeterol/ipratropium

	Per cartridge or blister
Salmeterol Xinafoate	72.6 microgram
Ipratropium Bromide	160 microgram
Lactose Ph. Eur.	to 12.5mg or to 25.0mg

5 Example 13: 50/320 salmeterol/ipratropium

	Per cartridge or blister
Salmeterol Xinafoate	72.6 microgram
Ipratropium Bromide	320 microgram
Lactose Ph. Eur.	to 12.5mg or to 25.0mg

Example 14: 50/80 salmeterol/ipratropium

	Per cartridge or blister
Salmeterol Xinafoate	72.5 microgram
Ipratropium Bromide	80 microgram
Lactose Ph. Eur.	to 12.5mg

10

The active ingredients were micronised and bulk blended with the lactose in the proportions given above (total blend size 4 kg). The blend was filled into specifically constructed double foil blister packs to be administered by a Diskus inhaler (Trademark of Glaxo Group Limited).

15

A similar method was used for the formulation of Example 15. The total blend size was 4 kg.

Example 15: 50/160 salmeterol/ipratropium

5

	Per cartridge or blister
Salmeterol Xinafoate	72.5 microgram
Ipratropium Bromide	160 microgram
Lactose Ph. Eur.	to 12.5mg

Claims

1. A pharmaceutical formulation comprising salmeterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and ipratropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients.
2. A pharmaceutical formulation comprising salmeterol xinafoate and ipratropium bromide, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients.
3. A pharmaceutical formulation according to claim 1 or 2 which is suitable for administration by inhalation.
4. A pharmaceutical formulation according to any of claims 1 to 3 which is a dry powder.
5. A pharmaceutical formulation according to any of claims 1 to 4 wherein the pharmaceutically acceptable carrier or excipient is lactose.
6. A pharmaceutical formulation according to any of claims 1 to 3 which is an aerosol formulation.
7. A pharmaceutical formulation according to any of claims 1 to 3 wherein the pharmaceutically acceptable carrier or excipient comprises a propellant.
8. A pharmaceutical formulation according to claim 6 wherein the propellant comprises 1,1,1,2-tetrafluoroethane and/or 1,1,1,2,3,3,3-heptafluoropropane.
9. A pharmaceutical formulation consisting essentially of particulate salmeterol or a pharmaceutically acceptable salt, solvate, or

- 5 physiologically functional derivative thereof and particulate ipratropium or
a pharmaceutically acceptable salt, solvate, or physiologically functional
derivative thereof, and optionally one or more other therapeutic
ingredients, and 1,1,1,2-tetrafluoroethane, 1,1,1,2,3,3,3-
heptafluoropropane or mixtures thereof as propellant.
- 10 10. A pharmaceutical formulation consisting essentially of particulate
salmeterol xinafoate and particulate ipratropium bromide, and optionally
one or more other therapeutic ingredients, and 1,1,1,2-tetrafluoroethane,
1,1,1,2,3,3,3-heptafluoropropane or mixtures thereof as propellant.
- 15 11. A method for the prophylaxis or treatment of a clinical condition in a
mammal, such as a human, for which a selective β_2 -adrenoreceptor
agonist and/or anticholinergic agent is indicated, which comprises
administration of a therapeutically effective amount of a pharmaceutical
formulation according to any one of claims 1 to 10.
- 20 12. A method according to claim 11 wherein the clinical condition is a
disease associated with reversible airways obstruction such as asthma,
chronic obstructive pulmonary disease (COPD), respiratory tract
infection or upper respiratory tract disease.
13. A method according to claim 11 wherein the clinical condition is chronic
obstructive pulmonary disease (COPD).

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 October 2001 (18.10.2001)

PCT

(10) International Publication Number
WO 01/76601 A3

(51) International Patent Classification⁷: **A61K 31/46**,
31/137 // (A61K 31/46, 31:137)

(74) Agent: **LEAROYD, Stephanie, Anne**; Glaxo Smith
Kline, Corporate Intellectual Property, Two New Horizons
Court, Brentford, Middlesex TW8 9EP (GB).

(21) International Application Number: **PCT/EP01/03889**

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(22) International Filing Date: **5 April 2001 (05.04.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
0008485.5 **7 April 2000 (07.04.2000)** **GB**

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **GLAXO
GROUP LIMITED [GB/GB]**; Glaxo Wellcome House,
Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).

Published:

with international search report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **JENKINS, Richard,
John [GB/US]**; SmithKline Beecham Corporation, Five
Moore Drive, Research Triangle Park, NC 27709 (US).
RICHARDS, David, Hugh [GB/GB]; GlaxoSmithKline,
Stockley Park West, Uxbridge, Middlesex UB11 1BT
(GB).

(88) Date of publication of the international search report:
21 February 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/76601 A3

(54) Title: **PHARMACEUTICAL COMPOSITIONS COMPRISING SALMETEROL AND IPRATROPIUM**

(57) Abstract: The present invention is concerned with pharmaceutical formulations comprising a combination of salmeterol and ipratropium and the use of such formulations in medicine, particularly in the prophylaxis and treatment of respiratory diseases.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/03889

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/46 A61K31/137 //(A61K31/46,31:137)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data, MEDLINE, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MATERA M G ET AL: "A COMBINATION WITH CLINICAL RECOMMENDED DOSAGES OF SALMETEROL AND IPRATROPIUM IS NOT MORE EFFECTIVE THAN SALMETEROL ALONE IN PATIENTSWITH CHRONIC OBSTRUCTIVE PULMONARY DISEASE" RESPIRATORY MEDICINE, BAILLIERE TINDALL, LONDON, GB, vol. 90, no. 8, 1996, pages 497-499, XP001021998 ISSN: 0954-6111 the whole document	1-13
X	WO 99 65464 A (BOEHRINGER INGELHEIM PHARMA) 23 December 1999 (1999-12-23) page 4, line 7-20; claims 1-20; example 1 -/--	1-13



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

25 October 2001

Date of mailing of the international search report

06/11/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Herrera, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/03889

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	<p>NOORD VAN J A ET AL: "LONG-TERM TREATMENT OF CHRONIC OBSTRUCTIVE PULMONARY DISEASE WITH SALMETEROL AND THE ADDITIVE EFFECT OF IPRATROPIUM"</p> <p>EUROPEAN RESPIRATORY JOURNAL, MUNKSGAARD INTERNATIONAL PUBLISHERS, COPENHAGEN, DK, vol. 15, no. 5, May 2000 (2000-05), pages 878-885, XP001021107</p> <p>ISSN: 0903-1936</p> <p>abstract</p> <p>-----</p>	1-13

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claims 11-13 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.1

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/03889

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9965464	A	23-12-1999	
		DE 19827178 A1	27-04-2000
		DE 19842963 A1	23-03-2000
		AU 4552199 A	05-01-2000
		BR 9911351 A	13-03-2001
		CN 1307470 T	08-08-2001
		EP 1087750 A1	04-04-2001
		NO 20006318 A	30-01-2001
		WO 9965464 A1	23-12-1999
